


Contents lists available at [SciVerse ScienceDirect](http://SciVerse.ScienceDirect.com)

Topology and its Applications

www.elsevier.com/locate/topol

Partial Hausdorff metric and Nadler's fixed point theorem on partial metric spaces

Hassen Aydi ^{a,*}, Mujahid Abbas ^b, Calogero Vetro ^c

^a Université de Sousse, Institut Supérieur d'Informatique et de Technologies de Communication de Hammam Sousse, Route GP1, Hammam Sousse-4011, Tunisia

^b Department of Mathematics, Lahore University of Management Sciences, 54792-Lahore, Pakistan

^c Dipartimento di Matematica e Informatica, Università degli Studi di Palermo, Via archirafi 34, 90123 Palermo, Italy

ARTICLE INFO

Article history:

Received 20 October 2011

Received in revised form 29 June 2012

Accepted 30 June 2012

MSC:

47H10

54H25

Keywords:

Partial Hausdorff metric

Multi-valued mappings

Nadler's fixed point theorem

ABSTRACT

In this paper, we introduce the concept of a partial Hausdorff metric. We initiate study of fixed point theory for multi-valued mappings on partial metric space using the partial Hausdorff metric and prove an analogous to the well-known Nadler's fixed point theorem. Moreover, we give a homotopy result as application of our main result.

© 2012 Elsevier B.V. All rights reserved.

1. Introduction and preliminaries

Let (X, d) be a metric space and $CB(X)$ denotes the collection of all nonempty closed and bounded subsets of X . For $A, B \in CB(X)$, define

$$H(A, B) := \max \left\{ \sup_{a \in A} d(a, B), \sup_{b \in B} d(b, A) \right\},$$

where $d(x, A) := \inf\{d(x, a) : a \in A\}$ is the distance of a point x to the set A . It is known that H is a metric on $CB(X)$, called the Hausdorff metric induced by the metric d .

Definition 1.1. Let X be any nonempty set. An element x in X is said to be a fixed point of a multi-valued mapping $T : X \rightarrow 2^X$ if $x \in Tx$, where 2^X denotes the collection of all nonempty subsets of X .

We recall that a multi-valued mapping $T : X \rightarrow CB(X)$ is said to be a contraction if

$$H(Tx, Ty) \leq kd(x, y)$$

for all $x, y \in X$ and for some k in $[0, 1)$.

The study of fixed points for multi-valued contractions using the Hausdorff metric was initiated by Nadler [14] who proved the following theorem.

* Corresponding author.

E-mail addresses: hassen.aydi@isima.rnu.tn (H. Aydi), mujahid@lums.edu.pk (M. Abbas), cvetro@math.unipa.it (C. Vetro).

Theorem 1.2. ([14]) Let (X, d) be a complete metric space and $T : X \rightarrow CB(X)$ be a contraction mapping. Then, there exists $x \in X$ such that $x \in Tx$.

Later, an interesting and rich fixed point theory was developed. The theory of multi-valued maps has application in control theory, convex optimization, differential equations and economics (see also [6]). On the other hand, Matthews [9] introduced the concept of a partial metric as a part of the study of denotational semantics of dataflow networks. He gave a modified version of the Banach contraction principle, more suitable in this context (see also [7,10]). In fact, (complete) partial metric spaces constitute a suitable framework to model several distinguished examples of the theory of computation and also to model metric spaces via domain theory (see, [5,8,9,11–13]). In this paper, we introduce the concept of a partial Hausdorff metric and extend the Nadler's fixed point theorem on partial metric spaces using the partial Hausdorff metric.

In the sequel the letters \mathbb{R} , \mathbb{R}^+ and \mathbb{N}^* will denote the set of all real numbers, the set of all nonnegative real numbers and the set of all positive integer numbers, respectively.

Consistent with [2,3,9], the following definitions and results will be needed in the sequel.

Definition 1.3. Let X be a nonempty set. A function $p : X \times X \rightarrow \mathbb{R}^+$ is said to be a partial metric on X if for any $x, y, z \in X$, the following conditions hold:

- (P₁) $p(x, x) = p(y, y) = p(x, y)$ if and only if $x = y$;
- (P₂) $p(x, x) \leq p(x, y)$;
- (P₃) $p(x, y) = p(y, x)$;
- (P₄) $p(x, z) \leq p(x, y) + p(y, z) - p(y, y)$.

The pair (X, p) is then called a partial metric space.

If $p(x, y) = 0$, then (P₁) and (P₂) imply that $x = y$. But the converse does not hold always.

A trivial example of a partial metric space is the pair (\mathbb{R}^+, p) , where $p : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is defined as $p(x, y) = \max\{x, y\}$ (see also [1]).

Example 1.4. ([9]) If $X = \{[a, b] : a, b \in \mathbb{R}, a \leq b\}$, then $p([a, b], [c, d]) = \max\{b, d\} - \min\{a, c\}$ defines a partial metric p on X .

For some more examples of partial metric spaces, we refer to [2,4,11,13].

Each partial metric p on X generates a T_0 topology τ_p on X which has as a base the family open p -balls $\{B_p(x, \varepsilon) : x \in X, \varepsilon > 0\}$, where $B_p(x, \varepsilon) = \{y \in X : p(x, y) < p(x, x) + \varepsilon\}$, for all $x \in X$ and $\varepsilon > 0$.

Observe (see [9, p. 187]) that a sequence $\{x_n\}$ in a partial metric space (X, p) converges to a point $x \in X$, with respect to τ_p , if and only if $p(x, x) = \lim_{n \rightarrow \infty} p(x, x_n)$.

If p is a partial metric on X , then the function $p^s : X \times X \rightarrow \mathbb{R}^+$ given by $p^s(x, y) = 2p(x, y) - p(x, x) - p(y, y)$, defines a metric on X .

Furthermore, a sequence $\{x_n\}$ converges in (X, p^s) to a point $x \in X$ if and only if

$$\lim_{n, m \rightarrow \infty} p(x_n, x_m) = \lim_{n \rightarrow \infty} p(x_n, x) = p(x, x). \quad (1.1)$$

Definition 1.5. ([9]) Let (X, p) be a partial metric space.

- (a) A sequence $\{x_n\}$ in X is said to be a Cauchy sequence if $\lim_{n, m \rightarrow \infty} p(x_n, x_m)$ exists and is finite.
- (b) (X, p) is said to be complete if every Cauchy sequence $\{x_n\}$ in X converges with respect to τ_p to a point $x \in X$ such that $\lim_{n \rightarrow \infty} p(x, x_n) = p(x, x)$. In this case, we say that the partial metric p is complete.

Lemma 1.6. ([2,9]) Let (X, p) be a partial metric space. Then:

- (a) A sequence $\{x_n\}$ in X is a Cauchy sequence in (X, p) if and only if it is a Cauchy sequence in metric space (X, p^s) .
- (b) A partial metric space (X, p) is complete if and only if the metric space (X, p^s) is complete.

2. Partial Hausdorff metric

Let (X, p) be a partial metric space. Let $CB^p(X)$ be the family of all nonempty, closed and bounded subsets of the partial metric space (X, p) , induced by the partial metric p . Note that Closedness is taken from (X, τ_p) (τ_p is the topology induced by p) and boundedness is given as follows: A is a bounded subset in (X, p) if there exist $x_0 \in X$ and $M \geq 0$ such that for all $a \in A$, we have $a \in B_p(x_0, M)$, that is, $p(x_0, a) < p(a, a) + M$.

For $A, B \in CB^p(X)$ and $x \in X$, define

$$p(x, A) = \inf\{p(x, a), a \in A\}, \quad \delta_p(A, B) = \sup\{p(a, B) : a \in A\} \quad \text{and} \\ \delta_p(B, A) = \sup\{p(b, A) : b \in B\}.$$

It is immediate to check that $p(x, A) = 0 \Rightarrow p^s(x, A) = 0$ where $p^s(x, A) = \inf\{p^s(x, a), a \in A\}$.

Remark 2.1. ([2]) Let (X, p) be a partial metric space and A any nonempty set in (X, p) , then

$$a \in \bar{A} \quad \text{if and only if} \quad p(a, A) = p(a, a), \quad (2.1)$$

where \bar{A} denotes the closure of A with respect to the partial metric p . Note that A is closed in (X, p) if and only if $A = \bar{A}$.

Now, we shall study some properties of mapping $\delta_p : CB^p(X) \times CB^p(X) \rightarrow [0, \infty)$.

Proposition 2.2. Let (X, p) be a partial metric space. For any $A, B, C \in CB^p(X)$, we have the following:

- (i) $\delta_p(A, A) = \sup\{p(a, a) : a \in A\}$;
- (ii) $\delta_p(A, A) \leq \delta_p(A, B)$;
- (iii) $\delta_p(A, B) = 0$ implies that $A \subseteq B$;
- (iv) $\delta_p(A, B) \leq \delta_p(A, C) + \delta_p(C, B) - \inf_{c \in C} p(c, c)$.

Proof. (i) From (2.1), if $A \in CB^p(X)$, then for all $a \in A$, we have $p(a, A) = p(a, a)$ as $\bar{A} = A$. Therefore $\delta_p(A, A) = \sup\{p(a, A) : a \in A\} = \sup\{p(a, a) : a \in A\}$.

(ii) Let $a \in A$. Since $p(a, a) \leq p(a, b)$ for all $b \in B$, therefore we have $p(a, a) \leq p(a, B) \leq \delta_p(A, B)$. From (i), we conclude that $\delta_p(A, A) = \sup\{p(a, a) : a \in A\} \leq \delta_p(A, B)$.

(iii) Suppose that $\delta_p(A, B) = 0$. Consequently $p(a, B) = 0$ for all $a \in A$. From (i) and (ii) it follows that $p(a, a) \leq \delta_p(A, B) = 0$ for all $a \in A$. That is, $p(a, a) = 0$ for all $a \in A$, and hence $p(a, B) = p(a, a)$ for all $a \in A$. Using (2.1), we have $a \in \bar{B} = B$ whenever $a \in A$ so $A \subseteq B$.

(iv) Let $a \in A$, $b \in B$ and $c \in C$. As

$$p(a, b) \leq p(a, c) + p(c, b) - p(c, c),$$

so we have

$$p(a, B) \leq p(a, c) + p(c, B) - p(c, c),$$

and

$$p(a, B) + p(c, c) \leq p(a, c) + \delta_p(C, B).$$

Since c is an arbitrary element of C , therefore we have

$$p(a, B) + \inf_{c \in C} p(c, c) \leq p(a, C) + \delta_p(C, B).$$

As a is an arbitrary element of A , so

$$\delta_p(A, B) \leq \delta_p(A, C) + \delta_p(C, B) - \inf_{c \in C} p(c, c). \quad \square$$

Let (X, p) be a partial metric space. For $A, B \in CB^p(X)$, define

$$H_p(A, B) = \max\{\delta_p(A, B), \delta_p(B, A)\}.$$

Proposition 2.3. Let (X, p) be a partial metric space. For all $A, B, C \in CB^p(X)$, we have

- (h1) $H_p(A, A) \leq H_p(A, B)$;
- (h2) $H_p(A, B) = H_p(B, A)$;
- (h3) $H_p(A, B) \leq H_p(A, C) + H_p(C, B) - \inf_{c \in C} p(c, c)$.

Proof. From (ii) of Proposition 2.2, we have $H_p(A, A) = \delta_p(A, A) \leq \delta_p(A, B) \leq H_p(A, B)$. By definition, (h2) holds obviously. Now using property (iv) of Proposition 2.2. We have

$$\begin{aligned} H_p(A, B) &= \max\{\delta_p(A, B), \delta_p(B, A)\} \\ &\leq \max\left\{\delta_p(A, C) + \delta_p(C, B) - \inf_{c \in C} p(c, c), \delta_p(B, C) + \delta_p(C, A) - \inf_{c \in C} p(c, c)\right\} \\ &= \max\{\delta_p(A, C) + \delta_p(C, B), \delta_p(B, C) + \delta_p(C, A)\} - \inf_{c \in C} p(c, c) \\ &\leq \max\{\delta_p(A, C), \delta_p(C, A)\} + \max\{\delta_p(C, B), \delta_p(B, C)\} - \inf_{c \in C} p(c, c) \\ &= H_p(A, C) + H_p(C, B) - \inf_{c \in C} p(c, c). \quad \square \end{aligned}$$

Corollary 2.4. Let (X, p) be a partial metric space. For $A, B \in CB^p(X)$ the following holds

$$H_p(A, B) = 0 \text{ implies that } A = B.$$

Proof. Let $H_p(A, B) = 0$. By definition of H_p , $\delta_p(A, B) = \delta_p(B, A) = 0$. Using (iii) of Proposition 2.2, we obtain that $A \subseteq B$ and $B \subseteq A$. Thus, $A = B$. \square

Remark 2.5. The converse of Corollary 2.4 is not true in general as it is clear from the following example.

Example 2.6. Let $X = [0, 1]$ be endowed with the partial metric $p : X \times X \rightarrow \mathbb{R}^+$ defined by

$$p(x, y) = \max\{x, y\}.$$

From (i) of Proposition 2.2, we have

$$H_p(X, X) = \delta_p(X, X) = \sup\{x : 0 \leq x \leq 1\} = 1 \neq 0.$$

In view of Proposition 2.3 and Corollary 2.4, we call the mapping $H_p : CB^p(X) \times CB^p(X) \rightarrow [0, +\infty)$, a partial Hausdorff metric induced by p .

Remark 2.7. It is easy to show that any Hausdorff metric is a partial Hausdorff metric. The converse is not true (see Example 2.6).

3. Fixed point of multi-valued contraction mapping

We start with the following lemma needed to prove our main result.

Lemma 3.1. Let (X, p) be a partial metric space, $A, B \in CB^p(X)$ and $h > 1$. For any $a \in A$, there exists $b = b(a) \in B$ such that

$$p(a, b) \leq hH_p(A, B). \quad (3.1)$$

Proof. If $A = B$, then from (i) of Proposition 2.2, we have

$$H_p(A, B) = H_p(A, A) = \delta_p(A, A) = \sup_{x \in A} p(x, x).$$

Let $a \in A$. Since $h > 1$, therefore we have

$$p(a, a) \leq \sup_{x \in A} p(x, x) = H_p(A, B) \leq hH_p(A, B).$$

Consequently $b = a$ satisfies (3.1). If $A \neq B$, suppose that there exists $a \in A$ such that $p(a, b) > hH_p(A, B)$ for all $b \in B$. This implies that $\inf\{p(a, y) : y \in B\} \geq hH_p(A, B)$. That is, $p(a, B) \geq hH_p(A, B)$. Note that

$$H_p(A, B) \geq \delta_p(A, B) = \sup_{x \in A} p(x, B) \geq p(a, B) \geq hH_p(A, B).$$

Since $A \neq B$, from Corollary 2.4 we have $H_p(A, B) \neq 0$ and above inequality gives $h \leq 1$, a contradiction. \square

Now, we state and prove our main result.

Theorem 3.2. Let (X, p) be a complete partial metric space. If $T : X \rightarrow CB^p(X)$ is a multi-valued mapping such that for all $x, y \in X$, we have

$$H_p(Tx, Ty) \leq kp(x, y) \quad (3.2)$$

where $k \in (0, 1)$. Then T has a fixed point.

Proof. Let $x_0 \in X$ and $x_1 \in Tx_0$. From Lemma 3.1 with $h = \frac{1}{\sqrt{k}}$, there exists $x_2 \in Tx_1$ such that $p(x_1, x_2) \leq \frac{1}{\sqrt{k}} H_p(Tx_0, Tx_1)$. As, $H_p(Tx_0, Tx_1) \leq kp(x_0, x_1)$ so $p(x_1, x_2) \leq \sqrt{k}p(x_0, x_1)$. For $x_2 \in Tx_1$, there exists $x_3 \in Tx_2$ such that $p(x_2, x_3) \leq \frac{1}{\sqrt{k}} H_p(Tx_1, Tx_2) \leq \sqrt{k}p(x_1, x_2)$. Continuing this process, we obtain a sequence $\{x_n\}$ in X such that

$$x_{n+1} \in Tx_n \quad \text{and} \quad p(x_{n+1}, x_n) \leq \sqrt{k}p(x_n, x_{n-1}) \quad \text{for all } n \geq 1. \quad (3.3)$$

Now from (3.3) and by the mathematical induction, we obtain

$$p(x_{n+1}, x_n) \leq (\sqrt{k})^n p(x_0, x_1) \quad \text{for all } n \in \mathbb{N}. \quad (3.4)$$

Using (3.4) and the property (P_4) of a partial metric, for any $m \in \mathbb{N}^*$, we have

$$\begin{aligned} p(x_n, x_{n+m}) &\leq p(x_n, x_{n+1}) + p(x_{n+1}, x_{n+2}) + \cdots + p(x_{n+m-1}, x_{n+m}) \\ &\leq (\sqrt{k})^n p(x_0, x_1) + (\sqrt{k})^{n+1} p(x_0, x_1) + \cdots + (\sqrt{k})^{n+m-1} p(x_0, x_1) \\ &= ((\sqrt{k})^n + (\sqrt{k})^{n+1} + \cdots + (\sqrt{k})^{n+m-1}) p(x_0, x_1) \\ &\leq \frac{(\sqrt{k})^n}{1 - \sqrt{k}} p(x_0, x_1) \rightarrow 0 \quad \text{as } n \rightarrow +\infty \quad (\text{since } 0 < k < 1). \end{aligned}$$

By the definition of p^s , we get for any $m \in \mathbb{N}^*$,

$$p^s(x_n, x_{n+m}) \leq 2p(x_n, x_{n+m}) \rightarrow 0 \quad \text{as } n \rightarrow +\infty. \quad (3.5)$$

This yields that $\{x_n\}$ is a Cauchy sequence in (X, p^s) . Since (X, p) is complete, then from Lemma 1.6, (X, p^s) is a complete metric space. Therefore, the sequence $\{x_n\}$ converges to some $x^* \in X$ with respect to the metric p^s , that is, $\lim_{n \rightarrow +\infty} p^s(x_n, x^*) = 0$. Again, from (1.1), we have

$$p(x^*, x^*) = \lim_{n \rightarrow +\infty} p(x_n, x^*) = \lim_{n \rightarrow +\infty} p(x_n, x_n) = 0. \quad (3.6)$$

Since $H_p(Tx_n, Tx^*) \leq kp(x_n, x^*)$, therefore

$$\lim_{n \rightarrow +\infty} H_p(Tx_n, Tx^*) = 0. \quad (3.7)$$

Now $x_{n+1} \in Tx_n$ gives that

$$p(x_{n+1}, Tx^*) \leq \delta_p(Tx_n, Tx^*) \leq H_p(Tx_n, Tx^*).$$

From (3.7), we get

$$\lim_{n \rightarrow +\infty} p(x_{n+1}, Tx^*) = 0. \quad (3.8)$$

On the other hand, we have

$$p(x^*, Tx^*) \leq p(x^*, x_{n+1}) + p(x_{n+1}, Tx^*).$$

Taking limit as $n \rightarrow +\infty$ and using (3.6) and (3.8), we obtain $p(x^*, Tx^*) = 0$. Therefore, from (3.6) ($p(x^*, x^*) = 0$), we obtain

$$p(x^*, x^*) = p(x^*, Tx^*),$$

which from (2.1) implies that $x^* \in \overline{Tx^*} = Tx^*$. \square

To underline the usefulness of partial metric, we give the following very simple illustrative examples.

Example 3.3. Let $X = \{0, 1, 4\}$ be endowed with the partial metric $p : X \times X \rightarrow \mathbb{R}^+$ defined by

$$p(x, y) = \frac{1}{4}|x - y| + \frac{1}{2} \max\{x, y\} \quad \text{for all } x, y \in X.$$

Note that $p(1, 1) = \frac{1}{2} \neq 0$ and $p(4, 4) = 2 \neq 0$, so p is not a metric on X . As $p^s(x, y) = |x - y|$ so (X, p) is a complete partial metric space.

Note that $\{0\}$ and $\{0, 1\}$ are bounded sets in (X, p) . In fact, if $x \in \{0, 1, 4\}$, then

$$\begin{aligned} x \in \overline{\{0\}} &\Leftrightarrow p(x, \{0\}) = p(x, x) \\ &\Leftrightarrow \frac{3}{4}x = \frac{1}{2}x \Leftrightarrow x = 0 \\ &\Leftrightarrow x \in \{0\}. \end{aligned}$$

Hence $\{0\}$ is closed with respect to the partial metric p . Also

$$\begin{aligned} x \in \overline{\{0, 1\}} &\Leftrightarrow p(x, \{0, 1\}) = p(x, x) \\ &\Leftrightarrow \min\left\{\frac{3}{4}x, \frac{1}{4}|x - 1| + \frac{1}{2} \max\{x, 1\}\right\} = \frac{1}{2}x \\ &\Leftrightarrow x \in \{0, 1\}. \end{aligned}$$

Hence $\{0, 1\}$ is closed with respect to the partial metric p .

Now, define the mapping $T : X \rightarrow CB^p(X)$ by

$$T(0) = T(1) = \{0\} \quad \text{and} \quad T(4) = \{0, 1\}.$$

We shall show that, for all $x, y \in X$, the contractive condition (3.2) is satisfied with $k = \frac{1}{2}$. For this, we consider the following cases:

- $x, y \in \{0, 1\}$. We have

$$H_p(T(x), T(y)) = H_p(\{0\}, \{0\}) = 0,$$

and (3.2) is satisfied obviously.

- $x \in \{0, 1\}$, $y = 4$. We have

$$\begin{aligned} H_p(T(0), T(4)) &= H_p(T(1), T(4)) \\ &= H_p(\{0\}, \{0, 1\}) \\ &= \max\{p(0, \{0, 1\}), \max\{p(0, 0), p(1, 0)\}\} \\ &= \frac{3}{4} \leq \frac{11}{8} = kp(1, 4) < \frac{3}{2} = kp(0, 4). \end{aligned}$$

- $x = y = 4$. We have

$$\begin{aligned} H_p(T(4), T(4)) &= H_p(\{0, 1\}, \{0, 1\}) \\ &= \sup\{p(x, x) : x \in \{0, 1\}\} \\ &= \max\{p(0, 0), p(1, 1)\} \\ &= \frac{1}{2} \leq 1 = kp(4, 4). \end{aligned}$$

Thus, all the hypotheses of Theorem 3.2 are satisfied. Here, $x = 0$ is a fixed point of T .

Example 3.4. Let $X = \{0, 1, 2\}$ be endowed with the partial metric $p : X \times X \rightarrow \mathbb{R}^+$ defined by

$$\begin{aligned} p(0, 0) &= p(1, 1) = 0, & p(2, 2) &= \frac{1}{4}, \\ p(0, 1) &= p(1, 0) = \frac{1}{3}, \\ p(0, 2) &= p(2, 0) = \frac{11}{24}, \\ p(1, 2) &= p(2, 1) = \frac{1}{2}. \end{aligned}$$

Define the mapping $T : X \rightarrow CB^p(X)$ by

$$T(0) = T(1) = \{0\} \quad \text{and} \quad T(2) = \{0, 1\}.$$

Note that, Tx is closed and bounded for all $x \in X$ under the given partial metric space (X, p) . We shall show that, for all $x, y \in X$, the contractive condition (3.2) is satisfied with $k = \frac{3}{4}$. For this, we consider the following cases:

- $x, y \in \{0, 1\}$. We have

$$H_p(T(x), T(y)) = H_p(\{0\}, \{0\}) = 0,$$

and (3.2) is satisfied obviously.

- $x \in \{0, 1\}$, $y = 2$. We have

$$\begin{aligned} H_p(T(1), T(2)) &= H_p(T(0), T(2)) \\ &= H_p(\{0\}, \{0, 1\}) \\ &= \max\{p(0, \{0, 1\}), \max\{p(0, 0), (1, 0)\}\} \\ &= \frac{1}{3} \leq \frac{11}{24}k = kp(0, 2) < \frac{1}{2}k = kp(1, 2). \end{aligned}$$

- $x = y = 2$. We have

$$\begin{aligned} H_p(T(2), T(2)) &= H_p(\{0, 1\}, \{0, 1\}) \\ &= \sup\{p(x, x) : x \in \{0, 1\}\} \\ &= \max\{p(0, 0), p(1, 1)\} \\ &= 0 < \frac{1}{4}k = kp(2, 2). \end{aligned}$$

Thus, all the conditions of Theorem 3.2 are satisfied. Here, $x = 0$ is a fixed point of T .

On the other hand, the metric p^s induced by the partial metric p is given by

$$\begin{aligned} p^s(0, 0) &= p^s(1, 1) = p^s(2, 2) = 0, \\ p^s(0, 1) &= p^s(1, 0) = p^s(0, 2) = p^s(2, 0) = \frac{2}{3}, \\ p^s(2, 1) &= p^s(1, 2) = \frac{3}{4}. \end{aligned}$$

Now, it is easy to show that Theorem 1.2 is not applicable in this case. Indeed, for $x = 0$ and $y = 2$, we have

$$\begin{aligned} H(T(0), T(2)) &= H(\{0\}, \{0, 1\}) \\ &= \max\{\sup\{p^s(0, \{0, 1\})\}, \sup\{p^s(\{0, 1\}, \{0\})\}\} \\ &= \max\left\{0, \frac{2}{3}\right\} = \frac{2}{3} \not\leq \frac{2}{3}k = kp^s(0, 2), \end{aligned}$$

for any $k \in (0, 1)$.

4. An application

In this section, as application of our main result, we derive a homotopy result.

Theorem 4.1. Let (X, p) be a complete partial metric space, A be an open subset of X and C be a closed subset of X , with $A \subset C$. Let $F : C \times [0, 1] \rightarrow CB^p(X)$ be an operator such that the following conditions are satisfied:

- $x \notin F(x, t)$ for each $x \in C \setminus A$ and each $t \in [0, 1]$,
- there exists $k \in (0, 1)$ such that for each $t \in [0, 1]$ and each $x, y \in C$ we have

$$H_p(F(x, t), F(y, t)) \leq kp(x, y),$$

(c) there exists a continuous function $\eta : [0, 1] \rightarrow \mathbb{R}$ such that

$$H_p(F(x, t), F(x, s)) \leq k|\eta(t) - \eta(s)| \quad \text{for all } t, s \in [0, 1] \text{ and each } x \in C,$$

(d) if $x \in F(x, t)$ then $F(x, t) = \{x\}$.

Then $F(\cdot, 0)$ has a fixed point if and only if $F(\cdot, 1)$ has a fixed point.

Proof. Consider the set

$$Q := \{t \in [0, 1] \mid x \in F(x, t) \text{ for some } x \in A\}.$$

Since $F(\cdot, 0)$ has a fixed point and (a) holds, we have that $0 \in Q$, so Q is a nonempty set. We will show that Q is both closed and open in $[0, 1]$, and so by the connectedness of $[0, 1]$ we are finished since $Q = [0, 1]$.

First, let us prove that Q is open in $[0, 1]$. Let $t_0 \in Q$ and $x_0 \in A$ with $x_0 \in F(x_0, t_0)$. Since A is open in (X, p) , there exists $r > 0$ such that $B_p(x_0, r) \subseteq A$. Consider $\varepsilon = r + p(x_0, x_0) - k(r + p(x_0, x_0)) > 0$. Since η is continuous on t_0 , there exists $\alpha(\varepsilon) > 0$ such that $|\eta(t) - \eta(t_0)| < \varepsilon$ for all $t \in (t_0 - \alpha(\varepsilon), t_0 + \alpha(\varepsilon))$.

Let $t \in (t_0 - \alpha(\varepsilon), t_0 + \alpha(\varepsilon))$, for $x \in \overline{B_p(x_0, r)} = \{x \in X \mid p(x_0, x) \leq p(x_0, x_0) + r\}$, we have

$$\begin{aligned} p(F(x, t), x_0) &\leq H_p(F(x, t), F(x_0, t_0)) \\ &\leq H_p(F(x, t), F(x, t_0)) + H_p(F(x, t_0), F(x_0, t_0)) \\ &\leq k|\eta(t) - \eta(t_0)| + kp(x, x_0) \\ &\leq k\varepsilon + k(p(x_0, x_0) + r) \\ &= k[r + p(x_0, x_0) - k(p(x_0, x_0) + r)] + k(p(x_0, x_0) + r) \\ &< r + p(x_0, x_0) - k(p(x_0, x_0) + r) + k(p(x_0, x_0) + r) \\ &= r + p(x_0, x_0). \end{aligned}$$

Then, for each fixed $t \in (t_0 - \alpha(\varepsilon), t_0 + \alpha(\varepsilon))$, $F(\cdot, t) : \overline{B_p(x_0, r)} \rightarrow CB^p(X)$ satisfies all the hypotheses of Theorem 3.2 and so $F(\cdot, t)$ has a fixed point in $\overline{B_p(x_0, r)} \subset C$. But this fixed point must be in A since (a) holds. Hence $(t_0 - \alpha(\varepsilon), t_0 + \alpha(\varepsilon)) \subseteq Q$ and therefore Q is open in $[0, 1]$.

Next, we show that Q is closed in $[0, 1]$. To see this, let $\{t_n\}_{n \in \mathbb{N}^*}$ be a sequence in Q with $t_n \rightarrow t^* \in [0, 1]$ as $n \rightarrow +\infty$. We must show that $t^* \in Q$. By the definition of Q , for all $n \in \mathbb{N}^*$, there exists $x_n \in A$ with $x_n \in F(x_n, t_n)$. Then, for $m, n \in \mathbb{N}^*$, we have using (d) and (h3)

$$\begin{aligned} p(x_n, x_m) &= H_p(F(x_n, t_n), F(x_m, t_m)) \\ &\leq H_p(F(x_n, t_n), F(x_n, t_m)) + H_p(F(x_n, t_m), F(x_m, t_m)) \\ &\leq k|\eta(t_n) - \eta(t_m)| + kp(x_n, x_m). \end{aligned}$$

This implies that

$$p(x_n, x_m) - kp(x_n, x_m) \leq k|\eta(t_n) - \eta(t_m)|,$$

and we get

$$p(x_n, x_m) \leq \frac{k}{1-k} |\eta(t_n) - \eta(t_m)|.$$

Since η is continuous and $\{t_n\}_{n \in \mathbb{N}^*}$ is convergent, letting $n, m \rightarrow +\infty$ in the above inequality, we obtain $\lim_{n, m \rightarrow +\infty} p(x_n, x_m) = 0$, that is, $\{x_n\}_{n \in \mathbb{N}^*}$ is a Cauchy sequence in (X, p) . Since (X, p) is complete, there exists $x^* \in C$ with $p(x^*, x^*) = \lim_{n \rightarrow +\infty} p(x^*, x_n) = \lim_{n, m \rightarrow +\infty} p(x_n, x_m) = 0$.

On the other hand, we have

$$\begin{aligned} p(x_n, F(x^*, t^*)) &\leq H_p(F(x_n, t_n), F(x^*, t^*)) \\ &\leq H_p(F(x_n, t_n), F(x_n, t^*)) + H_p(F(x_n, t^*), F(x^*, t^*)) \\ &\leq k|\eta(t_n) - \eta(t^*)| + kp(x_n, x^*). \end{aligned}$$

Letting $n \rightarrow +\infty$ in the above inequality, we get $\lim_{n \rightarrow +\infty} p(x_n, F(x^*, t^*)) = 0$ and so

$$p(x^*, F(x^*, t^*)) = \lim_{n \rightarrow +\infty} p(x_n, F(x^*, t^*)) = 0,$$

which implies that $x^* \in F(x^*, t^*)$, and since (a) holds, we have $x^* \in A$. Thus $t^* \in Q$ and Q is closed in $[0, 1]$.

One can use the same strategy to prove the reverse implication. This completes the proof. \square

Acknowledgements

The authors are grateful to the editor and referees for their valuable suggestions and critical remarks for improving this paper. The first author thanks Professor Bessem Samet for deep observations and fruitful discussions. The third author is supported by Università degli Studi di Palermo, Local University Project R. S. ex 60%.

Authors' contributions: All authors contributed equally and significantly in writing this paper. All authors read and approved the final manuscript.

References

- [1] M. Abbas, T. Nazir, Fixed point of generalized weakly contractive mappings in ordered partial metric spaces, *Fixed Point Theory Appl.* 2012 (1) (2012) 19 pp.
- [2] I. Altun, H. Simsek, Some fixed point theorems on dualistic partial metric spaces, *J. Adv. Math. Stud.* 1 (2008) 1–8.
- [3] I. Altun, F. Sola, H. Simsek, Generalized contractions on partial metric spaces, *Topology Appl.* 157 (2010) 2778–2785.
- [4] M.A. Bukatin, S.Yu. Shorina, Partial metrics and co-continuous valuations, in: M. Nivat, et al. (Eds.), *Foundations of Software Science and Computation Structure*, in: *Lecture Notes in Comput. Sci.*, vol. 1378, Springer, 1998, pp. 125–139.
- [5] L.j. Ćirić, B. Samet, H. Aydi, C. Vetro, Common fixed points of generalized contractions on partial metric spaces and an application, *Appl. Math. Comput.* 218 (2011) 2398–2406.
- [6] B. Damjanovic, B. Samet, C. Vetro, Common fixed point theorems for multi-valued maps, *Acta Math. Sci. Ser. B Engl. Ed.* 32 (2012) 818–824.
- [7] C. Di Bari, P. Vetro, Fixed points for weak φ -contractions on partial metric spaces, *Int. J. Eng., Contemp. Math. Sci.* 1 (1) (2011) 5–13.
- [8] R. Heckmann, Approximation of metric spaces by partial metric spaces, *Appl. Categ. Structures* 7 (1999) 71–83.
- [9] S.G. Matthews, Partial metric topology, in: *Proc. 8th Summer Conference on General Topology and Applications*, in: *Ann. New York Acad. Sci.*, vol. 728, 1994, pp. 183–197.
- [10] D. Paesano, P. Vetro, Suzuki's type characterizations of completeness for partial metric spaces and fixed points for partially ordered metric spaces, *Topology Appl.* 159 (3) (2012) 911–920.
- [11] S. Romaguera, A Kirk type characterization of completeness for partial metric spaces, *Fixed Point Theory Appl.* 2010 (2010), Article ID 493298, 6 pp.
- [12] S. Romaguera, O. Valero, A quantitative computational model for complete partial metric spaces via formal balls, *Math. Structures Comput. Sci.* 19 (2009) 541–563.
- [13] M.P. Schellekens, The correspondence between partial metrics and semivaluations, *Theoret. Comput. Sci.* 315 (2004) 135–149.
- [14] S.B. Nadler, Multivalued contraction mappings, *Pacific J. Math.* 30 (1969) 475–488.